

# Clinical Policy: Heart-Lung Transplant

Reference Number: CP.MP.132

Date of Last Revision: 02/25

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

## Description

Heart-lung transplantation is the treatment of choice for patients with both end-stage heart and end-stage lung disease. This policy establishes the medical necessity requirements for heart-lung transplants.

## Policy/Criteria

- I. It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that heart-lung transplant is **medically necessary** for members/enrollees who meet all of the following criteria:
  - A. End-stage heart and end-stage lung disease due to one of the following:
    1. Age  $\geq$  18 years and any of the following:
      - a. Irreversible primary pulmonary hypertension with severe heart failure;
      - b. Nonspecific Idiopathic severe pulmonary fibrosis;
      - c. Eisenmenger syndrome with irreversible pulmonary hypertension and heart failure;
      - d. Cystic fibrosis with severe heart failure;
      - e. Chronic obstructive pulmonary disease with severe heart failure;
      - f. Emphysema with severe heart failure;
      - g. Pulmonary fibrosis with uncontrollable pulmonary hypertension or heart failure;
      - h. Congenital heart disease (CHD) meeting one of the following:
        - i. Member/enrollee with single ventricle CHD and a Fontan circulation (total cardiopulmonary anastomosis) and one of the following:
          - a) Symptomatic heart failure (HF) and reduced systolic function (Class 1);
          - b) Symptomatic HF, preserved systolic function, and abnormal systemic ventricular filling pressures (Class 1);
          - c) Lymphatic abnormalities including plastic bronchitis and protein-losing enteropathy refractory to lymphatic interventions and medical management (Class 2a);
          - d) Cirrhosis or CKD attributed to chronically elevated central venous pressures (Class 2a);
        - ii. Member/enrollee with single ventricle CHD and one of the following:
          - a) Palliation to a shunted circulation or a superior cavo-pulmonary anastomosis (first procedure of a staged Fontan) and prohibitive risk for further single ventricle palliation;
          - b) Cyanotic heart disease with severe atrio-ventricular valve regurgitation and prohibitive risk for operative repair;
          - c) Pulmonary atresia with an intact ventricular septum, right ventricular dependent coronary circulation, and atresia of at least one aorto-coronary ostium;
        - iii. HF symptoms or ventricular arrhythmias refractory to medical, interventional, and device therapies (Class 1);
        - iv. Reactive pulmonary hypertension and a potential risk of developing fixed, irreversible elevation of PVR that could preclude heart failure in the future (Class 1);
      - i. Severe coronary artery disease or cardiomyopathy with irreversible pulmonary

**CLINICAL POLICY**  
**Heart-Lung Transplant**

- hypertension;
- j. Right ventricular failure with objective evidence of right ventricular fibrosis or infarction or refractory left ventricular failure;
- 2. Age < 18 years and any of the following:
  - a. Eisenmenger syndrome;
  - b. Heart re-transplant;
  - c. Lung re-transplant;
  - d. Pulmonary alveolar proteinosis;
  - e. Primary pulmonary hypertension;
  - f. Pulmonary vascular disease;
  - g. Restrictive cardiomyopathy;
  - h. Congenital heart disease meeting one of the following:
    - i. Member/enrollee with single ventricle CHD and a Fontan circulation (total cardiopulmonary anastomosis) and one of the following:
      - a) Symptomatic HF and reduced systolic function (Class 1);
      - b) Symptomatic HF, preserved systolic function, and abnormal systemic ventricular filling pressures (Class 1);
      - c) Lymphatic abnormalities including plastic bronchitis and protein-losing enteropathy refractory to lymphatic interventions and medical management (Class 2a);
      - d) Cirrhosis or CKD attributed to chronically elevated central venous pressures (Class 2a);
    - ii. Member/enrollee with single ventricle CHD and one of the following:
      - a) Palliation to a shunted circulation or a superior cavo-pulmonary anastomosis (first procedure of a staged Fontan) and prohibitive risk for further single ventricle palliation;
      - b) Cyanotic heart disease with severe atrio-ventricular valve regurgitation and prohibitive risk for operative repair;
      - c) Pulmonary atresia with an intact ventricular septum, right ventricular dependent coronary circulation, and atresia of at least one aorto-coronary ostium;
      - d) Neonatal hypoplastic left heart syndrome with high-risk features including HF symptoms, ventricular dysfunction, left ventricular-coronary artery fistulae;
    - iii. HF symptoms or ventricular arrhythmias refractory to medical, interventional, and device therapies (Class 1);
    - iv. Reactive pulmonary hypertension and a potential risk of developing fixed, irreversible elevation of PVR that could preclude heart failure in the future (Class 1);
    - v. Neonatal cyanotic CHD with high-risk features (Class 2a);
  - i. Cystic fibrosis with progressive, irreversible cardiac dysfunction;
  - j. Dilated cardiomyopathy;
- B. Meets the following disease severity criteria:
  - 1. Meets one of the following staging criteria:
    - a. Age ≥ 18 years: New York Heart Association classification of heart failure III or IV (Table 1);
    - b. Age < 18 years: American Heart Association Stage C or Stage D heart disease (Table 2);
  - 2. Life expectancy in the absence of cardiopulmonary disease ≥ two years;
- C. Does not have any of the following contraindications:
  - 1. HIV infection with detectable viral load unless all of the following are noted:
    - a. CD4 cell count > 200 cells/mm<sup>3</sup> for at least 3 months before transplantation;
    - b. Absence of active AIDS-defining opportunistic infection (unless treated efficaciously or

- prevented, can be included on the heart transplant waiting list) or malignancy;
- c. Member/enrollee is currently on effective antiretroviral therapy (ART);
- d. Member/enrollee does not have chronic wasting or severe malnutrition;
- 2. Inability to adhere to the regimen necessary to preserve the transplant, even with caregiver support;
- 3. Severe hypoplasia of the central branch pulmonary arteries or pulmonary veins;
- 4. Current episode of ongoing acute allograft rejection, even in the presence of graft vasculopathy, and retransplantation is requested;
- 5. Less than six months have passed since the primary transplantation, and retransplantation is requested;
- 6. Malignancy with high risk of recurrence or death related to cancer;
- 7. Acute renal failure with rising creatinine or on dialysis and low likelihood of recovery;
- 8. Acute liver failure or cirrhosis with portal hypertension or synthetic dysfunction, unless being considered for multi-organ transplant;
- 9. Stroke, acute coronary syndrome, or myocardial infarction (excluding demand ischemia) within 30 days;
- 10. Glomerular filtration rate < 30 mL/min/1.73m<sup>2</sup>, unless being considered for multi- organ transplant;
- 11. Septic shock;
- 12. Active extrapulmonary or disseminated infection;
- 13. Active tuberculosis infection;
- 14. Progressive cognitive impairment;
- 15. Other severe, uncontrolled medical condition expected to limit survival after transplant;
- 16. Active substance use or dependence (including current tobacco use, vaping, marijuana use [unless prescribed by a licensed practitioner] or intravenous drug use) without convincing evidence of risk reduction behaviors (unless urgent transplant timelines are present, in which case a commitment to reducing behaviors is acceptable). Serial blood and urine testing may be used to verify abstinence from substances that are of concern;
- 17. History of nicotine, tobacco, alcohol, or illicit drug use, without documentation noting abstinence from all (including nicotine replacement therapy) for ≥ six months prior to transplant;
- 18. Lung transplantation alone will restore right ventricular function.

**Table 1.** New York Heart Association (NYHA) Classifications of Heart Failure<sup>1</sup>

Classification	Characteristics
<b>Class I</b>	Patients with cardiac disease but without the resulting limitations in physical activity. Ordinary activity does not cause undue fatigue, palpitation, dyspnea, or anginal pain.
<b>Class II</b>	Patients with heart disease resulting in slight limitations of physical activity. They are comfortable at rest. Ordinary physical activity results in fatigue, palpitation, dyspnea or anginal pain.
<b>Class III</b>	Patients with cardiac disease resulting in marked limitation of physical activity. They are comfortable at rest. Less than ordinary physical activity causes fatigue, palpitation, dyspnea, or anginal pain.
<b>Class IV</b>	Patients with cardiac disease resulting in inability to carry on any physical activity without discomfort. The symptoms of cardiac insufficiency or of the anginal syndrome may be present even at rest. If any physical activity is undertaken, discomfort increases.

**Table 2.** American Heart Association (AHA) Heart Failure Stages<sup>1</sup>

**CLINICAL POLICY**  
**Heart-Lung Transplant**

Classification	Characteristics
<b>A</b>	Patients at high risk for heart failure but does not yet have symptoms or structural or functional heart disease.
<b>B</b>	Patients with no current or previous symptoms of heart failure but have structural heart disease, increased filling pressures in the heart, or other risk factors.
<b>C</b>	Patients with structural heart disease with current or previous symptoms of heart failure.
<b>D</b>	Patients who have heart failure with symptoms that interfere with daily life functions or result in recurrent hospitalizations despite continued guideline-directed medical therapy.

\*Note: Heart lung transplantations may be considered medically necessary for other congenital cardiopulmonary anomalies as determined upon individual case review.

**Background**

Heart-lung transplantation is a strong surgical option for selected patients with simultaneous end-stage heart failure and end-stage lung disease. Complex congenital heart disease with Eisenmenger syndrome is the most common indication for heart-lung transplantation, with other common indications to include primary pulmonary hypertension and cystic fibrosis. The frequency of heart-lung transplantation is limited due to the number of suitable donors, while the need for heart-lung transplantation has declined due to the availability of new medical therapies.<sup>2</sup>

Contraindications for combined heart-lung transplantation are similar to those for isolated heart and lung transplantation.<sup>2</sup> The International Society for Heart Lung Transplantation (ISHLT) provides listing criteria and best practice recommendations for heart transplants and for lung transplants.<sup>3,4,5</sup>

According to the 2019 ISHLT registry report, survival rates in adult patients who underwent heart-lung transplantation has steadily improved with an overall median survival rate of 3.7 years from 1992 through 2001 to 6.5 years from 2010 through 2017. This is comparable to primary lung transplantation but is inferior to the median survival rate of heart transplantation alone.<sup>2</sup>

**Coding Implications**

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2024, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
------------	-------------

**CLINICAL POLICY**  
**Heart-Lung Transplant**

33930	Donor cardiectomy-pneumonectomy (including cold preservation)
33933	Backbench standard preparation of cadaver donor heart/lung allograft prior to transplantation, including dissection of allograft from surrounding soft tissues to prepare aorta, superior vena cava, inferior vena cava, and trachea for implantation
33935	Heart-lung transplant with recipient cardiectomy-pneumonectomy

HCPCS Codes	Description
S2152	Solid organ(s), complete or segmental, single organ or combination of organs; deceased or living donor(s), procurement, transplantation, and related complications; including: drugs; supplies; hospitalization with outpatient follow-up; medical/surgical, diagnostic, emergency, and rehabilitative services, and the number of days of pre- and posttransplant care in the global definition

Reviews, Revisions, and Approvals	Revision Date	Approval Date
New policy.	06/17	06/17
References reviewed and updated. Replaced all instances of “member” with “member/enrollee.”	04/21	04/21
In B.2., removed “adequate functional status with the ability for rehabilitation.” Replaced contraindications of “history of psychological, behavioral, or cognitive disorders, poor family support structures, or documented noncompliance with previous therapies that could interfere with successful performance of care regimens after transplantation” and “current non-adherence to medical therapy...” with “Inability to adhere to the regimen necessary to preserve the transplant, even with caregiver support.” Changed “Review Date” in policy header to “Date of Last Revision,” and “Date” in the revision log header to “Revision Date.”	08/21	08/21
Annual review. References reviewed, updated, and reformatted. Updated I.C. with some contraindications from ISHLT 2021 guidelines. Background updated with no clinical significance.	02/22	02/22
Added specific congenital heart disease criteria to 2.i. Removed contraindication regarding specific congenital heart disease lesion.	05/22	05/22
Annual review completed. Removed pediatric indication of Alpha- 1 antitrypsin deficiency. Added “Lung transplantation alone will restore right ventricular function” to I.C. Updated I.C.10. to include “unless being considered for multi-organ transplant”. Criteria I.C.16. updated to exclude marijuana use when prescribed by a licensed practitioner and include required commitment to reducing substance use behaviors if urgent transplant timelines are present. ICD-10 diagnosis code table removed. Minor rewording with no clinical significance. References reviewed and updated. External specialists reviewed.	02/23	02/23

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Annual review. Added indication to criteria I.A.1.j. Expanded criteria I.C.1. to I.C.1.a. through c. Removed contraindication I.C.17., active peptic ulcer disease. References reviewed and updated.	02/24	02/24
Annual review. Changed I.A.1. to $\geq 18$ . I.A.1.a now reflects “severe” heart failure and I.A.1.b. now reflects “nonspecific idiopathic”. In I.A.1.h., “non-complex congenital...standard surgery” was removed, and now reflects “Congenital heart disease...”, adding I.A.1.h.i – iv., followed by i. and j. Age changed to $< 18$ in I.A.2 and in I.A.2.d, added “pulmonary” Changes made to I.A.2.h in addition to adding h.i.-v. Ages changed in I.B.1.a. “ $> 18$ ” and I.B.1.b. “ $\leq 18$ ”. In I.C.1.a, added “for at least ...transplantation”. Added I.C.1.d “Member/enrollee...severe malnutrition”. In I.C.8., added “unless being...transplant”. Reworded I.C.16.a. now reflecting “alcohol or illicit drug use”. Table 2 has been changed to the American Heart Association Heart Failure Stages, removing Heart Failure Stages in Pediatric Heart Disease. Background reviewed and updated. Coding verified. Internal and external specialist reviewed.	02/25	02/25

**References**

- American Heart Association Classes and Stages of Heart Failure. <https://www.heart.org/en/health-topics/heart-failure/what-is-heart-failure/classes-of-heart-failure>. Updated June 07, 2023. Accessed November 26, 2024.
- Singer LG, Mooney J. Heart-lung transplantation in adults. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated March 28, 2024. Accessed November 14, 2024.
- Mehra MR, Canter CE, Hannan MM, et al. The 2016 International Society for Heart Lung Transplantation listing criteria for heart transplantation: A 10-year update. *J Heart Lung Transplant*. 2016;35(1):1 to 23. doi:10.1016/j.healun.2015.10.023
- Leard LE, Holm AM, Valapour M, et al. Consensus document for the selection of lung transplant candidates: An update from the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. 2021;40(11):1349 to 1379. doi:10.1016/j.healun.2021.07.005
- Peled Y, Ducharme A, Kittleson M, et al. International Society for Heart and Lung Transplantation Guidelines for the Evaluation and Care of Cardiac Transplant Candidates-2024. *J Heart Lung Transplant*. 2024;43(10):1529-1628.e54. doi:10.1016/j.healun.2024.05.010
- Singh RK, Singh TP. Heart failure in children: Management. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated November 29, 2022. Accessed November 20, 2024.
- Helderman JH, Goral S. Gastrointestinal complications of transplant immunosuppression. *J Am Soc Nephrol*. 2002;13(1):277 to 287. doi:10.1681/ASN.V131277
- Vakil NM. Unusual causes of peptic ulcer disease. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated August 11, 2023. Accessed November 18, 2024.
- Weill D. Lung transplantation: indications and contraindications. *J Thorac Dis*. 2018;10(7):4574 to 4587. doi:10.21037/jtd.2018.06.141
- Spahr JE, West SC. Heart-lung transplantation: pediatric indications and outcomes. *J Thorac Dis*. 2014;6(8):1129 to 1137. doi:10.3978/j.issn.2072-1439.2014.07.05
- Le Pavec J, Hascoët S, Fadel E. Heart-lung transplantation: current indications, prognosis and specific considerations. *J Thorac Dis*. 2018;10(10):5946 to 5952.

**CLINICAL POLICY****Heart-Lung Transplant**

doi:10.21037/jtd.2018.09.115

12. Connolly HM, Frantz RP. Pulmonary hypertension in adults with congenital heart disease: disease-specific management. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated May 31, 2024. Accessed November 19, 2024.
13. Mancini D. Heart transplantation in adults: indications and contraindications. UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated October 25, 2024. Accessed November 18, 2024.
14. Hunt SA, Abraham WT, Chin MH, et al. 2009 focused update incorporated into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the International Society for Heart and Lung Transplantation [published correction appears in *Circulation*. 2010 Mar 30;121(12):e258]. *Circulation*. 2009;119(14):e391 to e479. doi:10.1161/CIRCULATIONAHA.109.192065
15. Pêgo-Fernandes PM. Heart-lung transplantation: a necessity. *J Bras Pneumol*. 2020;46(3):e20190273. Published 2020 Jun 5. doi:10.36416/1806-3756/e20190273
16. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;62(16):e147 to e239. doi:10.1016/j.jacc.2013.05.019
17. Stoller JK. Gene test interpretation: SERPINA1 (alpha-1 antitrypsin gene). UpToDate. [www.uptodate.com](http://www.uptodate.com). Updated March 25, 2024. Accessed November 19, 2024.
18. Westerdahl DE, Kobashigawa JA. Heart Transplantation for Advanced Heart Failure. *Cardiac Intensive Care*. 2018;;504-524.e2. <https://pubmed.ncbi.nlm.nih.gov/articles/PMC7161392/>. Accessed November 20, 2024.
19. Canter CE, Shaddy RE, Bernstein D, et al. Indications for heart transplantation in pediatric heart disease: a scientific statement from the American Heart Association Council on Cardiovascular Disease in the Young; the Councils on Clinical Cardiology, Cardiovascular Nursing, and Cardiovascular Surgery and Anesthesia; and the Quality of Care and Outcomes Research Interdisciplinary Working Group [published correction appears in *Circulation*. 2007 Apr 3;115(13):e385. Friedman, Allen H [corrected to Friedman, Alan H]]. *Circulation*. 2007;115(5):658 to 676. doi:10.1161/CIRCULATIONAHA.106.180449
20. Weill D, Benden C, Corris PA, et al. A consensus document for the selection of lung transplant candidates: 2014--an update from the Pulmonary Transplantation Council of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. 2015;34(1):1 to 15. doi:10.1016/j.healun.2014.06.014

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

## CLINICAL POLICY

### Heart-Lung Transplant

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

**Note: For Medicaid members/enrollees**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

**Note: For Medicare members/enrollees**, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.



## **CLINICAL POLICY**

### **Heart-Lung Transplant**

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporatio