

Clinical Policy: Leuprolide Acetate (Eligard, Fensolvi, Lupaneta Pack, Lupron Depot, Lupron Depot-Ped), Leuprolide mesylate (Camcevi)

Reference Number: CP.PHAR.173

Effective Date: 10.01.16

Last Review Date: 11.23

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Leuprolide acetate (Eligard[®], Fensolvi[®], Lupaneta Pack[®] [with norethindrone acetate tablets], Lupron Depot[®], Lupron Depot-Ped[®]) and leuprolide mesylate (Camcevi[™]) are gonadotropin-releasing hormone (GnRH) receptor agonists.

FDA Approved Indication(s)

Leuprolide acetate is indicated for:

- Palliative treatment of advanced prostate cancer:
 - Leuprolide acetate injection
- Treatment of advanced prostate cancer:
 - Lupron Depot (7.5, 22.5, 30, 45)
 - Eligard
- Initial management of the painful symptoms of endometriosis and for management of recurrence of symptoms:
 - Lupaneta Pack (3.75, 11.25)

Limitation(s) of use: Initial treatment course is limited to 6 months and use is not recommended longer than a total of 12 months due to concerns about adverse impact on bone mineral density.

- Management of endometriosis, including pain relief and reduction of endometriotic lesions; In combination with a norethindrone acetate for initial management of the painful symptoms of endometriosis and for management of recurrence of symptoms:

- Lupron Depot (3.75, 11.25)

Limitation(s) of use: total duration of therapy plus add-back therapy should not exceed 12 months due to concerns about adverse impact on bone mineral density

- Concomitant use with iron therapy for preoperative hematologic improvement of women with anemia caused by uterine leiomyomata [fibroids] for whom three months of hormonal suppression is deemed necessary:

- Lupron Depot (3.75, 11.25)

Limitation of use: not indicated for combination use with norethindrone acetate add-back therapy for the preoperative hematologic improvement of women with anemia caused by heavy menstrual bleeding due to fibroids

- Treatment of children with central precocious puberty (CPP):
 - Fensolvi
 - Leuprolide acetate
 - Lupron Depot-Ped (7.5, 11.25, 15, 30, 45)

Camcevi is indicated for the treatment of adult patients with advanced prostate cancer.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that leuprolide acetate, Camcevi, Eligard, Fensolvi, Lupaneta Pack, Lupron Depot, and Lupron Depot-Ped are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Prostate Cancer (must meet all):

1. Diagnosis of prostate cancer;
2. Request is for one of the following (a, b, c, or d):
 - a. Leuprolide acetate injection;
 - b. Camcevi;
 - c. Eligard;
 - d. Lupron Depot;
3. Prescribed by or in consultation with an oncologist or urologist;
4. Age \geq 18 years;
5. Request meets one of the following (a, b, c, or d):*
 - a. Leuprolide acetate injection (SC): Dose does not exceed 1 mg per day;
 - b. Camcevi (SC): Dose does not exceed 42 mg per 6 months;
 - c. Eligard (SC)/Lupron Depot (IM): Dose does not exceed 7.5 mg per month, 22.5 mg per 3 months, 30 mg per 4 months, 45 mg per 6 months;
 - d. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member's renewal date, whichever is longer

B. Endometriosis (must meet all):

1. Diagnosis of endometriosis;
2. Request is for one of the following (a or b):
 - a. Lupron Depot (3.75 mg, 11.25 mg);
 - b. Lupaneta Pack (3.75 mg, 11.25 mg);
3. Prescribed by or in consultation with a gynecologist;
4. Age \geq 18 years;
5. Endometriosis as a cause of pain is one of the following (a or b):
 - a. Surgically confirmed;
 - b. Both of the following (i and ii):
 - i. Clinically suspected;
 - ii. Failure of a 3-month trial of one of the following within the last year, unless clinically adverse effects are experienced or all are contraindicated (1, 2, or 3):
 - 1) A nonsteroidal anti-inflammatory drug (*see Appendix B for examples*);

- 2) An oral or injectable depot contraceptive (*see Appendix B for examples*);
- 3) A progestin (*see Appendix B for examples*);
6. For members currently receiving treatment with leuprolide, total duration of therapy has not exceeded 12 months;
7. Dose does not exceed 3.75 mg per month or 11.25 mg per 3 months.

Approval duration:

Medicaid/HIM – 6 months

Commercial – 6 months or to member’s renewal date, whichever is longer

C. Uterine Fibroids (must meet all):

1. Diagnosis of anemia secondary to uterine leiomyomata (fibroids);
2. Diagnosis is confirmed by ultrasound;
3. Request is for Lupron Depot (3.75 mg, 11.25 mg);
4. Prescribed by or in consultation with gynecologist;
5. Age \geq 18 years;
6. Lupron Depot is prescribed concurrently with iron therapy;
7. Prescribed preoperatively to reduce fibroid size and improve hematologic control;
8. For members currently receiving treatment with leuprolide, total duration of therapy has not exceeded 3 months per treatment course;
9. Dose does not exceed 3.75 mg per month or 11.25 mg per 3 months.

Approval duration: 3 months

D. Central Precocious Puberty (must meet all):

1. Member meets one of the following (a or b):
 - a. Diagnosis of CPP confirmed by all of the following (i, ii, and iii):
 - i. Elevated basal luteinizing hormone (LH) level $> 0.2 - 0.3$ mIU/mL (dependent on type of assay used) and/or elevated leuprolide-stimulated LH level $> 3.3 - 5$ IU/L (dependent on type of assay used);
 - ii. Difference between bone age and chronological age was > 1 year (bone age-chronological age);
 - iii. Age at onset of secondary sex characteristics (1 or 2):
 - 1) Female: < 8 years;
 - 2) Male: < 9 years;
 - b. Request is for diagnostic use;
2. Request is for one of the following (a, b, or c):
 - a. Fensolvi;
 - b. Leuprolide acetate;
 - c. Lupron Depot-Ped: 7.5 mg, 11.25 mg, 15 mg, 30 mg, 45 mg;
3. Prescribed by or in consultation with a pediatric endocrinologist;
4. Member meets one of the following age requirements (a or b):
 - a. Female: 2 - 11 years;
 - b. Male: 2 - 12 years;
5. Dose does not exceed the following (a, b, c, or d):
 - a. Diagnostic use: Leuprolide acetate: 20 mcg/kg or as needed;
 - b. Therapeutic use: Fensolvi: 45 mg per 6 months;

- c. Therapeutic use: Leuprolide acetate (SC): Initial: 50 mcg/kg per day; titrate dose upward by 10 mcg/kg per day if down-regulation is not achieved (higher mg/kg doses may be required in younger children).
- d. Therapeutic use: Lupron Depot-Ped (IM): 15 mg per month (1-month formulation), 30 mg per 3 months (3-month formulation) or 45 mg per 6 months (6-month formulation) (dosing is weight-based for a 1-month and a 3-month formulations).

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member’s renewal date, whichever is longer

E. Breast and Ovarian Cancer (off-label) (must meet all):

1. Diagnosis of hormone receptor-positive breast cancer or ovarian cancer (including fallopian tube and primary peritoneal cancer);
2. Request is for Lupron Depot;
3. Prescribed by or in consultation with an oncologist;
4. Age \geq 18 years;
5. Request meets one of the following (a, b, or c):*
 - a. Breast or ovarian cancer: Dose does not exceed 3.75 mg per month or 11.25 mg per 3 months;
 - b. Ovarian cancer: Dose does not exceed 7.5 mg per month, 11.25 mg per 3 months, 22.5 mg per 3 months;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member’s renewal date, whichever is longer

F. Gender Dysphoria, Gender Transition (off-label) (must meet all):

1. Diagnosis of gender dysphoria or request is for gender transition;
2. Request is for a leuprolide product other than Lupaneta Pack;
3. Prescribed by or in consultation with both of the following (a and b):
 - a. An endocrinologist;
 - b. A provider with expertise in gender dysphoria and transgender medicine based on a certified training program or affiliation with local transgender health services (e.g., mental health professional such as psychologist, psychiatrist, see *Appendix D*);
4. Age and pubertal development - meets one of the following (a or b):
 - a. Member is $<$ 18 years of age and has reached or passed through Tanner Stage 2*;
**Age ranges approximating Tanner Stage 2 pubertal development extend from 8 to 13 years of age in girls and 9 to 14 years of age in boys.*
 - b. Member is \geq 18 years of age and has failed to achieve physiologic hormone levels with gender-affirming hormonal therapy (e.g., estrogen, testosterone) unless contraindicated or clinically significant adverse effects are experienced;
5. Member demonstrates understanding of expected GnRH analogue treatment outcomes and has given consent for such treatment;

6. If member has a psychiatric comorbidity, member is followed by mental health provider;
7. Psychosocial support will be provided during treatment;
8. Dose is within FDA maximum limit for any FDA-approved indication (see Section V) or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member’s renewal date, whichever is longer

G. Salivary Gland Tumors (off-label) (must meet all):

1. Diagnosis of salivary gland tumors;
2. Disease is androgen receptor positive and recurrent, unresectable, or metastatic;
3. Prescribed by or in consultation with an oncologist;
4. Request is for one of the following (a or b):
 - a. Eligard;
 - b. Lupron Depot;
5. Dose is within FDA maximum limit for any FDA-approved indication (see Section V) or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member’s renewal date, whichever is longer

H. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Prostate Cancer (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving leuprolide acetate injection, Camcevi, Eligard, or

- Lupron Depot for prostate cancer and has received this medication for at least 30 days;
2. Request is for one of the following (a, b, c, or d):
 - a. Leuprolide acetate injection;
 - b. Camcevi;
 - c. Eligard;
 - d. Lupron Depot;
 3. Member is responding positively to therapy;
 4. If request is for a dose increase, request meets one of the following (a, b, c, or d):*
 - a. Leuprolide acetate injection (SC): New dose does not exceed 1 mg per day;
 - b. Camcevi (SC): New dose does not exceed 42 mg per 6 months;
 - c. Eligard (SC)/Lupron Depot (IM): New dose does not exceed 7.5 mg per month, 22.5 mg per 3 months, 30 mg per 4 months, 45 mg per 6 months;
 - d. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member's renewal date, whichever is longer

B. Endometriosis (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Request is for Lupron Depot/Lupaneta Pack (3.75 mg, 11.25 mg);
3. Member is responding positively to therapy as evidenced by, including but not limited to, improvement in any of the following parameters: improvement in dysmenorrhea, dyspareunia, pelvic pain/induration/tenderness, or size of endometrial lesions;
4. Total duration of leuprolide therapy has not exceeded 12 months;
5. If request is for a dose increase, new dose does not exceed 3.75 mg per month or 11.25 mg per 3 months.

Approval duration:

Medicaid/HIM – up to a total treatment duration of 12 months

Commercial – 6 months or to member's renewal date, whichever is longer

C. Uterine Fibroids:

1. Re-authorization is not permitted. Members must meet the initial approval criteria.

Approval duration: Not applicable

D. Central Precocious Puberty (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;

- b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Request is for one of the following (a, b, or c):
 - a. Fensolvi;
 - b. Leuprolide acetate;
 - c. Lupron Depot-Ped: 7.5 mg, 11.25 mg, 15 mg, 30 mg, 45 mg;
3. Member is responding positively to therapy as evidenced by, including but not limited to, improvement in any of the following parameters: decreased growth velocity, cessation of menses, softening of breast tissue or testes, arrested pubertal progression;
4. Member meets one of the following age requirements (a or b):
 - a. Female: ≤ 11 years;
 - b. Male: ≤ 12 years;
5. If request is for a dose increase, new dose does not exceed one of the following (a, b, or c):
 - a. Leuprolide acetate (SC): Initial: 50 mcg/kg per day; titrate dose upward by 10 mcg/kg per day if down-regulation is not achieved (higher mg/kg doses may be required in younger children);
 - b. Lupron Depot-Ped (IM): 15 mg per month (1-month formulation), 30 mg per 3 months (3-month formulation) or 45 mg per 6 months (6-month formulation) (dosing is weight-based for a 1-month and a 3-month formulations);
 - c. Fensolvi: 45 mg per 6 months.

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member’s renewal date, whichever is longer

E. Breast and Ovarian Cancer (off-label) (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Lupron Depot for hormone receptor-positive breast cancer or ovarian cancer and has received this medication for at least 30 days;
2. Request is for Lupron Depot;
3. Member is responding positively to therapy;
4. If request is for a dose increase, request meets one of the following (a, b, or c):*
 - a. Breast or ovarian cancer: New dose does not exceed 3.75 mg per month or 11.25 mg per 3 months;
 - b. Ovarian cancer: New dose does not exceed 7.5 mg per month, 11.25 mg per 3 months, 22.5 mg per 3 months;
 - c. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member’s renewal date, whichever is longer

F. Gender Dysphoria, Gender Transition (off-label) (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Request is for a leuprolide product other than Lupaneta Pack;
3. Member is responding positively to therapy (e.g., member continues to meet their individual goals of therapy for gender dysphoria);
4. If request is for a dose increase, new dose is within FDA maximum limit for any FDA-approved indication (see Section V) or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member’s renewal date, whichever is longer

G. Salivary Gland Tumors (off-label) (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Eligard or Lupron Depot for salivary gland tumors and has received this medication for at least 30 days ;
2. Request is for one of the following (a or b):
 - a. Eligard;
 - b. Lupron Depot;
3. Member is responding positively to therapy;
4. If request is for a dose increase, new dose is within FDA maximum limit for any FDA-approved indication (see Section V) or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or to member’s renewal date, whichever is longer

H. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CPP: central precocious puberty
 DSM-5: Diagnostic and Statistical Manual of Mental Disorders, 5th edition
 FDA: Food and Drug Administration
 GnRH: gonadotropin-releasing hormone

LH: luteinizing hormone
 NCCN: National Comprehensive Cancer Network
 WPATH: World Professional Association for Transgender Health

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
NSAIDs*: ibuprofen, naproxen, fenoprofen, ketoprofen, mefenamic acid, meclufenamate, indomethacin, tolmetin, diclofenac, etodolac, diflunisal, meloxicam, piroxicam	Endometriosis Varies – refer to specific prescribing information	Varies – refer to specific prescribing information
Combined oral estrogen-progesterone contraceptives: ethinyl estradiol + (desogestrel, ethynodiol diacetate, drospirenone, etonogestrel, levonorgestrel, norelgestromin, norethindrone, norgestimate, or norgestrel); estradiol valerate + dienogest; mestranol + norethindrone	Endometriosis 1 tablet PO QD (may vary per specific prescribing information)	1 tablet per day (may vary per specific prescribing information)
Progestin-only oral contraceptives: norethindrone	Endometriosis 0.35 mg PO QD	0.35 mg per day
Depot injection progestin contraceptives: medroxyprogesterone acetate	Endometriosis IM: 150 mg per 3 months (every 13 weeks)	See regimen

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	SC: 104 mg per 3 months (every 12 to 14 weeks)	

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

**Examples provided may not be all-inclusive*

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Known hypersensitivity to GnRH, GnRH agonist analogs or any of the components of the individual products (all leuprolide products);
 - Pregnancy (all leuprolide products except Camcevi, Eligard);
 - Lupron 3.75 mg/11.25 mg and Lupaneta Pack:
 - Undiagnosed abnormal vaginal bleeding;
 - Breast-feeding;
 - If used with norethindrone acetate:
 - Thrombophlebitis, thromboembolic disorders, cerebral apoplexy, or a past history of these conditions;
 - Markedly impaired liver function or liver disease;
 - Known or suspected carcinoma of the breast.
- Boxed warning(s): None reported

Appendix D: General Information

- World Professional Association for Transgender Health (WPATH) offers their Global Education Institute (GEI) Certified Training Courses: Best Practices in Transgender Medical and Mental Health Care. Additionally, the following link provides a search tool to locate WPATH member providers: <https://www.wpath.org/provider/search>
- Transgender Care Therapy Certification Training is also offered by the International Transgender Certification Association (ITCA). Professionals with expertise in transgender care can be located using the following search tool: <https://transgendercertification.com/locate-a-professional/>
- The WPATH Standards of Care Version 8 recommend that adolescents are managed by a multidisciplinary care team that involves both medical and mental health professionals. The list of key disciplines includes but is not limited to: adolescent medicine/primary care, endocrinology, psychology, psychiatry, speech/language pathology, fertility, social work, support staff, and the surgical team. The need to include a healthcare professional with some expertise in mental health does not dictate the inclusion of a psychologist, psychiatrist or social work in every assessment. Instead, a general practitioner, nurse or other qualified clinician could fulfill this requirement as long as they have sufficient expertise to diagnose gender incongruence, recognize mental health concerns, distinguish between these concerns and gender dysphoria, incongruence or diversity, assist a transgender person in care planning and preparing for gender affirmative medical and surgical treatments, and refer to a mental health professional if needed.

Appendix E: Additional Information on Diagnosis-specific HCPCS Codes, Billable Units, and Day Supply

Diagnosis	Requested Product	HCPCS Code	Billable Units	Day Supply
Prostate Cancer	Leuprolide acetate, per 1 mg	J9218	14	14
	Lupron Depot 1-Month & Eligard 7.5 mg	J9217	1	28
	Lupron Depot 3-Month & Eligard 22.5 mg		3	84
	Lupron Depot 4-Month & Eligard 30 mg		4	112
	Lupron Depot 6-Month & Eligard 45 mg		6	168
	Camcevi 6-Month 42 mg	NA	NA	168
Endometriosis, Uterine Fibroids	Lupron Depot 1-Month 3.75 mg	J1950	1	28
	Lupron Depot 3-Month 11.25 mg		3	84
Central Precocious Puberty	Leuprolide acetate, per 1 mg	J9218	14	14
	Lupron Depot-Ped 7.5 mg	J1950	2	28
	Lupron Depot-Ped 11.25 mg		3	28
	Lupron Depot-Ped 15 mg		4	28
	Lupron Depot-Ped 30 mg		8	84
	Lupron Depot-Ped 45 mg		12	168
	Fensolvi 45 mg kit		12	168
Breast Cancer	Lupron Depot 1-Month 3.75 mg	J1950	1	28
Ovarian Cancer	Lupron Depot 1-Month 3.75 mg	J1950	1	28
	Lupron Depot 3-Month 11.25 mg		3	84
Salivary Gland Tumors	Lupron Depot 1-Month & Eligard 7.5 mg	J9217	1	28
	Lupron Depot 3-Month & Eligard 22.5 mg		3	84

NA – not available

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Leuprolide acetate injection	Prostate cancer	Camcevi (SC) – 42 mg every 6 months	See regimen
Leuprolide acetate (Lupron Depot 7.5, 22.5, 30, 45)		Leuprolide acetate injection (SC): 1 mg per day	See regimen
		Lupron Depot (IM) - 7.5 mg per month; 22.5 mg per 3 months; 30 mg per 4 months; 45 mg per 6 months	See regimen
Leuprolide acetate (Eligard 7.5, 22.5, 30, 45)		Eligard (SC) - 7.5 mg per month; 22.5 mg per 3 months; 30 mg per 4 months; 45 mg per 6 months	See regimen
Leuprolide mesylate (Camcevi)			

Drug Name	Indication	Dosing Regimen	Maximum Dose
Leuprolide acetate (Lupron Depot 3.75, 11.25) Leuprolide acetate (Lupaneta Pack 3.75, 11.25)	Endometriosis	Lupron Depot/Lupaneta Pack (IM) - 3.75 mg per month; 11.25 mg per 3 months	See regimen
Leuprolide acetate (Lupron Depot 3.75)	Uterine fibroids	Lupron Depot (IM) - 3.75 mg/month, 11.25 mg per 3 months	See regimen
Leuprolide acetate injection Leuprolide acetate (Lupron Depot-Ped 7.5, 11.25, 15 [1 mo]; 11.25, 30 [3 mo]); 45 [6 mo] Fensolvi (leuprolide acetate)	CPP	Leuprolide acetate (SC): <ul style="list-style-type: none"> • Diagnostic: 20 mcg/kg or as needed; • Treatment: Initial: 50 mcg/kg/day; titrate dose upward by 10 mcg/kg/day if down-regulation is not achieved (higher mg/kg doses may be required in younger children). 	See regimen
		Lupron Depot-Ped (IM): Monthly administration weight-based starting dose: 7.5 mg (\leq 25 kg), 11.25 mg ($>$ 25 to 37.5 kg), 15 mg ($>$ 37.5 kg) (increase as needed to 15 mg/month); 3-month administration: 11.25 mg or 30 mg; 6-month administration: 45 mg	See regimen
		Fensolvi (SC): 45 mg once every six months	See regimen
Leuprolide acetate (Lupron Depot 3.75)	Breast cancer (off-label)	Lupron Depot (IM) 3.75 mg per month, 11.25 mg per 3 months	See regimen
Leuprolide acetate (Lupron Depot 3.75, 11.25)	Ovarian cancer (off-label)	Lupron Depot (IM) 3.75 mg per month, 11.25 mg per 3 months	See regimen
Leuprolide acetate (Lupron Depot 7.5, 22.5) Leuprolide acetate (Eligard 7.5, 22.5, 30, 45)	Salivary Gland tumors (off-label)	Lupron Depot (IM) - 7.5 mg per month; 22.5 mg per 3 months. Eligard (SC) - 7.5 mg per month; 22.5 mg per 3 months; 30 mg per 4 months; 45 mg per 6 months	See regimen

VI. Product Availability

Drug Name	Availability
Leuprolide acetate injection	Kit: 2.8 mL multi-dose vial (1 mg/0.2 mL)
Leuprolide acetate (Eligard)	Kit: 7.5 mg (1 month), 22.5 mg (3 month), 30 mg (4 month), 45 mg (6 month)
Leuprolide acetate and norethindrone tablets (Lupaneta Pack)	Pack: 3.75 mg leuprolide acetate syringe (1 month) with 5 mg norethindrone tablets Pack: 11.25 mg leuprolide acetate syringe (3 month) with 5 mg norethindrone tablets
Leuprolide acetate (Lupron Depot)	Prefilled syringe: 7.5 mg (1 month), 22.5 mg (3 month), 30 mg (4 month), 45 mg (6 month)
Leuprolide acetate (Lupron Depot 3.75)	Prefilled syringe: 3.75 mg (1 month)
Leuprolide acetate (Lupron Depot 11.25)	Prefilled syringe: 11.25 mg (3 month)
Leuprolide acetate (Lupron Depot-Ped)	Prefilled syringe: 7.5 mg (1 month), 11.25 mg (1 month), 15 mg (1 month) Prefilled syringe: 11.25 mg (3 month), 30 mg (3 month) Prefilled syringe: 45 mg (6 month)
Leuprolide acetate (Fensolvi)	Kit: syringe A: prefilled with diluent for reconstitution and syringe B: prefilled with 45 mg lyophilized leuprolide acetate powder
Leuprolide mesylate (Camcevi)	Injection emulsion: 42 mg

VII. References

1. Leuprolide Acetate Injection Prescribing Information. Princeton, NJ: Sandoz Inc.; August 2011. Available at https://www.accessdata.fda.gov/drugsatfda_docs/label/2011/074728s011lbl.pdf. Accessed July 12, 2023.
2. Camcevi Prescribing Information. Taipei City, Taiwan: Foresee Pharmaceuticals Co.; November 2022. Available at https://www.accordbiopharma.com/our-therapies/camcevi/camcevi_pi.pdf. Accessed July 12, 2023.
3. Eligard Prescribing Information. Fort Collins CO: TOLMAR Pharmaceuticals, Inc.; July 2023. Available at https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/021343s047s048,021379s049s050,021488s044s045,021731s044s047lbl.pdf. Accessed July 26, 2023.
4. Fensolvi Prescribing Information. Fort Collins, CO: Tolmar, Inc.; November 2022. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/213150s003lbl.pdf. Accessed July 12, 2023.
5. Lupaneta Pack 3.75 Prescribing Information. North Chicago, IL: AbbVie Inc.; December 2022. http://rxabbvie.com/pdf/lupaneta_3_75_pi.pdf. Accessed June 30, 2023.
6. Lupaneta Pack 11.25 Prescribing Information. North Chicago, IL: AbbVie Inc.; December 2022. http://rxabbvie.com/pdf/lupaneta_11_25_pi.pdf. Accessed June 30, 2023.
7. Lupron Depot 3.75 Prescribing Information. North Chicago, IL: AbbVie Inc.; January 2023 Available at https://www.rxabbvie.com/pdf/lupron3_75mg.pdf. Accessed June 30, 2023.

8. Lupron Depot 11.25 mg Prescribing Information. North Chicago, IL: AbbVie Inc.; March 2020. Available at https://www.rxabbvie.com/pdf/lupron3month11_25mg.pdf . Accessed June 30, 2023.
 9. Lupron Depot 7.5, 22.5, 30, 45 mg Prescribing Information. North Chicago, IL: AbbVie, Inc., April 2022. Available at http://www.rxabbvie.com/pdf/lupronuro_pi.pdf. Accessed June 30, 2023.
 10. Lupron Depot-PED Prescribing Information. North Chicago, IL: AbbVie Inc.; April 2023. Available at: <http://www.rxabbvie.com/pdf/lupronpediatric.pdf>. Accessed June 30, 2023.
 11. National Comprehensive Cancer Network Drugs and Biologics Compendium. Leuprolide acetate. Available at [nccn.org](http://www.nccn.org). Accessed July 12, 2023.
 12. National Comprehensive Cancer Network Drugs and Biologics Compendium. Leuprolide acetate for depot suspension. Available at [nccn.org](http://www.nccn.org). Accessed July 12, 2023.
 13. National Comprehensive Cancer Network Drugs and Biologics Compendium. Leuprolide mesylate. Available at [nccn.org](http://www.nccn.org). Accessed July 12, 2023.
 14. National Comprehensive Cancer Network. Prostate cancer (Version 1.2023). Available at https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf. Accessed July 12, 2023.
 15. National Comprehensive Cancer Network. Breast cancer (Version 4.2023). Available at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed July 12, 2023.
 16. National Comprehensive Cancer Network. Ovarian cancer (Version 2.2023). Available at https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf. Accessed July 12, 2023.
 17. National Comprehensive Cancer Network. Head and Neck Cancers (Version 2.2023). Available at https://www.nccn.org/professionals/physician_gls/pdf/head-and-neck.pdf . Accessed July 12, 2023.
 18. Committee on Practice Bulletins - Gynecology. Management of endometriosis. July 2010 (reaffirmed 2016); 116(1): 223-236.
 19. Donnez J, Dolmans MM. Uterine fibroids management: From the present to the future. Hum Reprod Update. November 2016; 22(6): 665-686.
 20. MS DeLaCruz, EM Buchanan. Uterine fibroids: Diagnosis and treatment. Am Fam Physician. January 15, 2017; 95(2): 100-107.
 21. American College of Obstetricians and Gynecologists' Committee on Practice Bulletins–Gynecology. Management of Symptomatic Uterine Leiomyomas: ACOG Practice Bulletin, Number 228. Obstet Gynecol. 2021 Jun 1;137(6):e100-e115
 22. Marret H, Fritel X, Ouldamer L et al. Therapeutic management of uterine fibroid tumors: Updated French guidelines. Eur J Obstet Gynecol Reprod Biol. December 2012; 165(2): 156–164.
 23. Kaplowitz P, Bloch C. Evaluation and referral of children with signs of early puberty. Pediatrics. 2016; 137(1): e20153732.
 24. Falcone T, Flyckt R. Clinical Management of Endometriosis. Obstetrics & Gynecology. 2018; 131(3):p557-571.
 25. Management of Symptomatic Uterine Leiomyomas: ACOG Practice Bulletin Number 228, Clinical Management Guidelines for Obstetrician-Gynecologists. Obstet Gynecol. 2021 Jun 1;137(6):e100-e115.
- Gender Dysphoria*
26. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th ed. Arlington, VA: American Psychiatric Association Publishing; 2013.

27. Hembree WC, Cohen-Kettenis PT, Gooren L, et al. Endocrine treatment of gender-dysphoric/gender-incongruent persons: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*, November 2017, 102(11):3869–3903.
28. Rafferty J. Ensuring comprehensive care and support for transgender and gender-diverse children and adolescents. Policy statement. *American Academy of Pediatrics. Pediatrics*; 142(4), October 2018:e20182162.
29. Deutsch MB. Guidelines for the primary and gender-affirming care of transgender and gender nonbinary people. Center of Excellence for Transgender Health. Department of Family & Community Medicine, University of California, San Francisco; 2nd Ed, published June 17, 2016.
30. Coleman E, Bockting W, Botzer M, et al. Standards of care for the health of transsexual, transgender, and gender nonconforming people. *WPATH: World Professional Association for Transgender Health*. 7th version; 2012. Available at https://www.wpath.org/media/cms/Documents/SOC%20v7/SOC%20V7_English2012.pdf?t=1613669341. Accessed July 12, 2023.
31. Wylie KR, Fung R, Boshier C, Rotchell M. Recommendations of endocrine treatment for patients with gender dysphoria. *Sexual and Relationship Therapy* Vol. 24, No. 2, May 2009, 175–187.
32. Emmanuel M, Bokor BR. *Tanner Stages*. Treasure Island, FL: StatPearls Publishing; 2019 Jan. Available at <https://www.ncbi.nlm.nih.gov/books/NBK470280/>. Last update: May 13, 2019. Accessed July 12, 2023.
33. Micromedex[®] Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed July 12, 2023.
34. WPATH: World Professional Association for Transgender Health Standards of Care Version 8 Draft. Available at: <https://www.wpath.org/soc8>. Accessed July 12, 2023.

Coding Implications*

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J1950	Injection, leuprolide acetate (for depot suspension), per 3.75 mg
J9217	Leuprolide acetate (for depot suspension), 7.5 mg
J9218	Leuprolide acetate, per 1 mg
J9219	Leuprolide acetate implant, 65 mg
J1951	Injection, leuprolide acetate for depot suspension (Fensolvi), 0.25 mg
J1952	Leuprolide injectable, Camcevi, 1 mg

*See Appendix E: Additional Information on Diagnosis-specific HCPCS Codes, Billable Units, and Day Supply

Reviews, Revisions, and Approvals	Date	P&T Approval Date
4Q 2019 annual review: added Commercial and HIM-Medical Benefit line of business, added notation that Lupron Depot-Ped (3	08.01.19	11.19

Reviews, Revisions, and Approvals	Date	P&T Approval Date
month) 11.25 mg strength is non-formulary for HIM; for prostate cancer added urologist specialist option; references reviewed and updated.		
3Q 2020 annual review: revised HIM-Medical Benefit to HIM line of business; added Fensolvi (new dosage form) to the policy for Central Precocious Puberty; added off-label NCCN indication and criteria for salivary gland tumor; references reviewed and updated.	05.07.20	08.20
Added Appendix D: Additional Information on Diagnosis-specific HCPCS Codes, Billable Units, and Day Supply.	01.25.21	
Clarified that diagnosis is not necessary for diagnostic use for CPP.	04.19.21	05.21
3Q 2021 annual review: for endometriosis and uterine fibroid indications added requirements for total duration of therapy per prescribing information; for uterine fibroids continuation of therapy revised to restrict re-authorization and require use of initial approval criteria as each preoperative treatment course would be evaluated individually; revised salivary gland tumor to allow continuity of care and revised initial approval duration from duration of request or through the end of contract year to 12 months to align with other oncology approval durations; for gender dysphoria continuation of therapy added requirement that request is not for Lupaneta Pack to align with initial approval criteria; for ovarian cancer added Lupron Depot 7.5 mg and 22.5 mg strengths per NCCN; references to HIM.PHAR.21 revised to HIM.PA.154; references reviewed and updated.	04.13.21	08.21
4Q 2021 annual review: RT4: Added Camcevi, a new dosage form of existing product [Lupron Depot] with same indication for prostate cancer; added gender transition to gender dysphoria criteria set; clarified breast cancer should be hormone receptor-positive; references reviewed and updated.	07.08.21	11.21
For gender dysphoria or request is for gender transition modified prescriber requirements to allow experts in transgender medicine based on a certified training program or affiliation with local transgender health services; modified Appendix D to E; for general information Appendix D added resources for transgender provider search tools and examples of training programs.	12.06.21	
4Q 2022 annual review: modified Commercial approval duration to 6 months or to member's renewal date, whichever is longer; added HCPCS codes for Fensolvi and Camcevi; for Lupron Depot (7.5, 22.5, 30, 45) updated FDA-approved indication to include non-palliative treatment of advanced prostate cancer; references reviewed and updated. Template changes applied to other diagnoses/indications and continued therapy section.	07.21.22	11.22

Reviews, Revisions, and Approvals	Date	P&T Approval Date
For oncology indications, removed references to specific Lupron Depot strengths.	01.05.23	
For breast cancer maximum dosing added option for 11.25 mg per 3 months.	03.21.23	
RT4: added new dosage form (45 mg) for 6-month dosing regimen for Lupron Depot-Ped	04.19.23	
4Q 2023 annual review: for uterine fibroids added requirement that Lupron Depot is prescribed concurrently with iron therapy per FDA indication and revised Commercial approval duration to 3 months as treatment per label is limited to three months; for gender dysphoria continuation of therapy added example of positive response to therapy; references reviewed and updated. RT4: updated Eligard FDA-approved indication per prescribing information for use in the treatment of advanced prostate cancer.	06.30.23	11.23
Corrected units for basal luteinizing hormone level from mIU/L to mIU/mL for CPP.	02.08.24	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan

retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.